



PATIENT INFORMATION (Complete or Fax Existing Chart) PRESCRIBER INFORMATION

Name: _____ DOB: _____ Address: _____ City, State, Zip: _____ Phone: _____ Alt. Phone: _____ Email: _____ SS#: _____ Gender: <input type="checkbox"/> M <input type="checkbox"/> F Weight: _____ (lbs) Ht: _____ Allergies: _____	Prescriber Name: _____ State License: _____ NPI #: _____ Tax ID: _____ Address: _____ City, State, Zip: _____ Phone: _____ Fax: _____ Office Contact: _____ Phone: _____
---	--

INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)

Primary Insurance: _____ Plan #: _____ Group #: _____ RX Card (PBM): _____ BIN: _____ PCN: _____	Secondary Insurance (If Applicable): _____ Plan #: _____ Group #: _____ RX Card (PBM): _____ BIN: _____ PCN: _____
--	--

CLINICAL INFORMATION

Primary ICD-10 Code (Please specify diagnosis): _____
 Secondary ICD-10 Code (Please specify diagnosis): _____
 Number of Gout Flare per year: _____ Glucose-6-phosphate dehydrogenase (G6PD) Testing (Please Provide Results)
 Serum Uric Acid Level at Baseline: _____ mg/dl Serum Uric Acid Level Prior to Infusion: _____ mg/dl
 Is Patient Currently Prescribed and/or Taking Immunomodulation (MTX)? Yes No
Past/Current Medical History (select all that apply)
 CHF BP: Controlled Uncontrolled Pregnant Breast feeding Anaphylactic reaction to previous IV therapy
 Tophus Joints affected: _____

KRYSTEXXA® ORDERS

Prescription type: New start Restart Continued therapy Total Doses Received: _____ Date of Last Injection/Infusion: _____

Medication	Dose/Frequency	Refills
<input type="checkbox"/> Krystexxa® (pegloticase)	<input type="checkbox"/> 8 mg IV every 2 weeks <input type="checkbox"/> Other: _____	_____
Pre-Medication	Dose/Strength	Directions
<input type="checkbox"/> Acetaminophen	<input type="checkbox"/> 500mg	<input type="checkbox"/> Take 1-2 tablets PO prior to infusion or post-infusion as directed
<input type="checkbox"/> Diphenhydramine	<input type="checkbox"/> 25mg IV/PO <input type="checkbox"/> 50mg IV/PO	<input type="checkbox"/> Take 1 tablet PO prior to infusion or as directed OR <input type="checkbox"/> Inject contents of 1 vial IV prior to infusion or as directed
<input type="checkbox"/> Methylprednisolone	<input type="checkbox"/> 40mg <input type="checkbox"/> 100mg <input type="checkbox"/> 125mg	<input type="checkbox"/> Inject contents of 1 vial IV prior to infusion or as directed <input type="checkbox"/> Other: Inject 100mg IV 30 minutes prior to infusion
<input type="checkbox"/> _____	_____	_____

INFUSION REACTION ORDERS

Mild reaction protocol:
 Diphenhydramine 25mg IV, one time, for pruritus.
If symptoms worsen, see orders for moderate to severe reactions.

Moderate reaction protocol:
 Acetaminophen 650mg PO, one time, for pyrexia or rigors
 Diphenhydramine 50mg IV, one time, for pruritus or urticaria
 Famotidine 20mg IV, one time, for, for pruritus or urticaria

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.



- Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms
- If symptoms worsen, see interventions for severe reactions*
- Severe reaction protocol: (Call 911 if initiated):**
- Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)
 - Diphenhydramine 50mg IV, one time, for respiratory symptoms, edema, or anaphylaxis
 - Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis
 - Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms
 - Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or worsen

FLUSHING & LOCKING ORDERS

Flushing Protocol (>66lbs/33kg)

<p>PIV and Midline:</p> <ul style="list-style-type: none"> <input checked="" type="checkbox"/> 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion 	<p>Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:</p> <ul style="list-style-type: none"> <input checked="" type="checkbox"/> 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw
---	--

Locking Protocol (>66lbs/33kg)

<p>PIV and Midline:</p> <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Heparin Sodium 10 units/mL 1mL IV final flush post normal saline flush 	<p>PICC:</p> <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush 	<p>Implanted Port, Tunneled Catheter, and Non-tunneled Catheter:</p> <ul style="list-style-type: none"> <input checked="" type="checkbox"/> Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush
---	--	---

**** May substitute Dextrose 5% in Water, or alternative, for 0.9% Sodium Chloride, when indicated due to incompatibility with medications being infused**

SIGNATURE

We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.

X _____
Prescriber Signature

Date: _____

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.