



| PATIENT INFORMATION (Complete or Fax Existing Chart) | | PRESCRIBER INFORMATION | |
|--|-------|--|--|
| Name: _____ DOB: _____ Address: _____ City, State, Zip: _____ Phone: _____ Alt. Phone: _____ Email: _____ SS#: _____ Gender: <input type="checkbox"/> M <input type="checkbox"/> F Weight: _____ (lbs) Ht: _____ Allergies: _____ | | Prescriber Name: _____ State License: _____ NPI #: _____ Tax ID: _____ Address: _____ City, State, Zip: _____ Phone: _____ Fax: _____ Office Contact: _____ Phone: _____ | |
| INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back) | | | |
| Primary Insurance: _____ Plan #: _____ Group #: _____ RX Card (PBM): _____ BIN: _____ PCN: _____ | | Secondary Insurance (If Applicable): _____ Plan #: _____ Group #: _____ RX Card (PBM): _____ BIN: _____ PCN: _____ | |
| CLINICAL INFORMATION | | | |
| <input type="checkbox"/> G70.00 Myasthenia gravis without (acute) exacerbation <input type="checkbox"/> G70.01: Myasthenia gravis with (acute) exacerbation <input type="checkbox"/> Other: _____ MG-ADL* score (if known): _____ Adverse reactions with previous MG treatments: _____ If so, what MG treatment caused the reaction: _____ | | | |
| ORDERS | | | |
| Prescription type: <input type="checkbox"/> New start <input type="checkbox"/> Restart <input type="checkbox"/> Continued therapy Total Doses Received: _____ Date of Last Injection/Infusion: _____ | | | |
| Medication | Route | Strength/Formulation | Directions |
| Imaavy™ | IV | 1200mg/6.5mL single-dose vial 300mg/ 1.62ml single-dose vial | <input type="checkbox"/> Loading dose at Week 0: Infuse 30mg/kg intravenously once over at least 30 minutes rounding to an easily measurable dose when clinically appropriate. <input type="checkbox"/> Maintenance dose: Infuse 15mg/kg intravenously over at least 15 minutes every 2 weeks thereafter rounding to an easily measurable dose when clinically appropriate. |
| Pre-Medication | | Dose/Strength | Directions |
| <input type="checkbox"/> Acetaminophen | | <input type="checkbox"/> 500mg | <input type="checkbox"/> Take 1-2 tablets PO prior to infusion or post-infusion as directed |
| <input type="checkbox"/> Diphenhydramine | | <input type="checkbox"/> 25mg IV/PO <input type="checkbox"/> 50mg IV/PO | <input type="checkbox"/> Take 1 tablet PO prior to infusion or as directed OR <input type="checkbox"/> Inject contents of 1 vial IV prior to infusion or as directed |
| <input type="checkbox"/> Methylprednisolone | | <input type="checkbox"/> 40mg <input type="checkbox"/> 100mg <input type="checkbox"/> 125mg | <input type="checkbox"/> Inject contents of 1 vial IV prior to infusion or as directed <input type="checkbox"/> Other: Inject 100mg IV 30 minutes prior to infusion |
| <input type="checkbox"/> _____ | | _____ | _____ |
| INFUSION REACTION ORDERS | | | |
| Mild reaction protocol: <input checked="" type="checkbox"/> Diphenhydramine 25mg IV, one time, for pruritus. <i>If symptoms worsen, see orders for moderate to severe reactions.</i> Moderate reaction protocol: <input checked="" type="checkbox"/> Acetaminophen 650mg PO, one time, for pyrexia or rigors <input checked="" type="checkbox"/> Diphenhydramine 50mg IV, one time, for pruritus or urticaria | | | |



☒ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms

If symptoms worsen, see interventions for severe reactions

Severe reaction protocol: (Call 911 if initiated):

☒ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)

☒ Diphenhydramine 50mg IV, one time, for respiratory symptoms, edema, or anaphylaxis

☒ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis

☒ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms

☒ Epinephrine 0.3mg/0.3mL IM into mid-antrolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or worsen

FLUSHING & LOCKING ORDERS

Flushing Protocol (>66lbs/33kg)

PIV and Midline:

☒ 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion

Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:

☒ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw

Locking Protocol (>66lbs/33kg)

PIV and Midline:

☒ Heparin Sodium 10 units/mL 1mL IV final flush post normal saline flush

PICC:

☒ Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush

Implanted Port, Tunneled Catheter, and Non-tunneled Catheter:

☒ Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush

**** May substitute Dextrose 5% in Water, or alternative, for 0.9% Sodium Chloride, when indicated due to incompatibility with medications being infused**

SIGNATURE

We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.

X _____

Prescriber Signature

Date: _____

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.

CONFIDENTIALITY STATEMENT: This facsimile and documents accompanying this transmission contain confidential health information that is legally privileged. This information is intended only for the use of the individual or entity named above. The authorized recipient of this information is prohibited from disclosing this information to any other party unless required to do so by law or regulation. If you are not the intended recipient, you are hereby notified that any disclosure, copying, distribution, or action taken in reliance on the contents of these documents is strictly prohibited. If you have received this information in error, please notify the sender at the address and telephone number set forth herein and arrange for return or destruction of the material. In no event should such material be read by anyone other than the named addressee, except by express authority of the sender to the named addressee.