TALIS HEALTHCARE
AN INFUSION MANAGEMENT COMPANY

LEQVIO®

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

PATIENT INFORMATION (Complete or Fax Existing Chart)

Name: _____ DOB: _____
Address: _____
City, State, Zip: _____
Phone: _____ Alt. Phone: _____
Email: _____ SS#: _____
Gender: ☐ M ☐ F Weight: _____(lbs) Ht: _____
Allergies: _____

PRESCRIBER INFORMATION

Prescriber Name: _____
State License: _____
NPI #: _____ Tax ID: _____
Address: _____
City, State, Zip: _____
Phone: _____ Fax: _____
Office Contact: _____ Phone: _____

INSURANCE INFORMATION – AND – Send a copy of the patient's prescription/insurance cards (front & back)

Primary Insurance: _____
Plan #: _____
Group #: _____
RX Card (PBM): _____
BIN: _____ PCN: _____

Secondary Insurance (If Applicable): _____
Plan #: _____
Group #: _____
RX Card (PBM): _____
BIN: _____ PCN: _____

CLINICAL INFORMATION

Primary diagnosis section (select one diagnosis code; complete ICD-10-CM to the highest level of specificity)

Hypercholesterolemia: ☐ E78.00 ☐ E78.2 ☐ E78.49 ☐ E78.5 Familial hypercholesterolemia (eg, HeFH): ☐ E78.011 ☐ E78.019

Other (specify ICD-10-CM): _____

Secondary diagnosis(es) (please complete if Hypercholesterolemia above is selected; complete ICD-10-CM to highest level of specificity)

Clinical ASCVD:

☐ I2 ___.____ Ischemic heart disease ☐ I6 ___.____ Cerebrovascular disease ☐ I70._____ Atherosclerosis ☐ I73._____ Other peripheral vascular disease

☐ Other (specify ICD-10-CM): _____

AND/OR

Other clinical risk factors: ☐ E11.____ Diabetes mellitus ☐ I10.____ Hypertension ☐ Other (specify ICD-10-CM): _____

LDL-C level: _____ Date taken: _____ (MM/DD/YYYY) Current LDL-C lowering treatment(s): _____

ORDERS

Prescription type: ☐ New start ☐ Restart ☐ Continued therapy Total Doses Received: _____ Date of Last Injection/Infusion: _____

Medication	Dose/Frequency	Qty/Refills
☐ Leqvio (Inclisiran)	☐ Induction: Administer 284mg subcutaneously at day 0, month 3 then every 6 months ☐ Maintenance: Administer 284mg subcutaneously every 6 months ☐ Other: _____	Quantity: _____ Refills: _____

Pre-Medication	Dose/Strength	Directions
☐ Acetaminophen	☐ 500mg	☐ Take 1-2 tablets PO prior to infusion or post-infusion as directed
☐ Diphenhydramine	☐ 25mg IV/PO ☐ 50mg IV/PO	☐ Take 1 tablet PO prior to infusion or as directed OR ☐ Inject contents of 1 vial IV prior to infusion or as directed
☐ Methylprednisolone	☐ 40mg ☐ 100mg ☐ 125mg	☐ Inject contents of 1 vial IV prior to infusion or as directed ☐ Other: Inject 100mg IV 30 minutes prior to infusion
☐ _____	_____	_____

INFUSION REACTION ORDERS

TALIS HEALTHCARE
AN INFUSION MANAGEMENT COMPANY

LEQVIO®

Please Fax Completed Form To: 888-898-9113

Please Send a Copy of The Patient's Insurance Cards (Front & Back)

Mild reaction protocol:

☒ Diphenhydramine 25mg IV, one time, for pruritus.

If symptoms worsen, see orders for moderate to severe reactions.

Moderate reaction protocol:

☒ Acetaminophen 650mg PO, one time, for pyrexia or rigors

☒ Diphenhydramine 50mg IV, one time, for pruritus or urticaria

☒ Famotidine 20mg IV, one time, for, for pruritus or urticaria

☒ Methylprednisolone 125mg IV, one time, for respiratory or neurologic symptoms

If symptoms worsen, see interventions for severe reactions

Severe reaction protocol: (Call 911 if initiated):

☒ Titrate oxygen via continuous flow per nasal cannula or face mask to maintain spO2 of greater than ninety-five percent (>95%)

☒ Diphenhydramine 50mg IV,one time, for respiratory symptoms, edema, or anaphylaxis

☒ Methylprednisolone 125mg IV, one time, for respiratory symptoms, edema, or anaphylaxis

☒ Sodium Chloride 0.9% 500mL IV over 30-60 min, one time, for cardiovascular symptoms

☒ Epinephrine 0.3mg/0.3mL IM into mis-anterolateral aspect of thigh of anaphylaxis, may repeat x1 in 5-15 minutes if symptoms are not resolved or worsen

FLUSHING & LOCKING ORDERS

Flushing Protocol (>66lbs/33kg)

PIV and Midline:	**Implanted Port, PICC, Tunneled Catheter, and Non-tunneled Catheter:**
☒ 0.9% Sodium Chloride 2-5mL IV flush before and after each infusion	☒ 0.9% Sodium Chloride 5mL IV flush before infusion/lab draw and 10mL IV flush after infusion/lab draw

Locking Protocol (>66lbs/33kg)

PIV and Midline:	**PICC:**	**Implanted Port, Tunneled Catheter, and Non-tunneled Catheter:**
☒ Heparin Sodium 10 units/mL 1mL IV final flush post normal saline flush	☒ Heparin Sodium 10 units/mL 3mL IV final flush post normal saline flush	☒ Heparin Sodium 100 units/mL 3-5mL IV final flush post normal saline flush

**** May substitute Dextrose 5% in Water, or alternative, for 0.9& Sodium Chloride, when indicated due to incompatibility with medications bring infused**

SIGNATURE

We hereby authorize Talis Healthcare LLC to provide all supplies and additional services (nursing/patient training) required to provide and deliver the medicine as prescribed in this referral.

X_____ Date: _____

Prescriber Signature

To ensure payment by insurance carrier, please include supporting clinical documentation for specified ICD 10 Code, demographic, and insurance information along with faxed order. Initial appointment will be verified upon insurance approval.